

PERMANENT MAKE-UP PRE-PROCEDURE INFORMATION

Be prepared for the color intensity of your procedure to be significantly sharper, brighter or darker than what is expected for the final outcome. It will take time for this transition, based on how quickly the outer layer of your skin exfoliates.

While the injected tones may initially simulate the exact color and tone desired, it will not always remain a perfect match. Injected tones are constant, while your own skin tones will vary depending on exposure to cold, heat, sun and circulatory changes. For example, if you tan your skin and had a scar camouflaged your surrounding skin will be darker in appearance than the treated area.

- Since delicate skin or sensitive areas may swell slightly or redden, some clients prefer to not make any social plans for a day or two following the procedure.
- Do not wear make-up on the day of your procedure. Do bring your lip or brow pencils with you. If you choose to wear make-up, please bring your make-up cleanser with you.
- Any tweezing or waxing should be done at least 48 hours prior to the the procedure; electrolysis or laser no less than five days before. Do not resume any method of hair removal for at least two weeks.
- Any eyelash or eyebrow tinting or eyelash curling should be done no sooner than 48 hours before or two weeks after the procedure.
- Do not wear contact lenses during or immediately following the eyeliner procedure. Remember to bring your glasses. You may resume wearing your contact lenses as soon as your eyes return to their pre-tattooed condition. Bring sunglasses on the day of your procedure since your eyes may be light sensitive post-procedure.
- Following the eyeliner procedure, as a safety precaution, we recommend that you have someone available to accompany you or drive you home.
- If you are having lip procedures and have any history of cold sores/fever blisters/herpes simplex, you will be required to take medication prior to the lip procedure.
- Refrain from the use of alcohol, aspirin, aspirin-containing medications, Coumadin, or other anti-inflammatory medications for 7 days before and 2 days after the procedure. However: **NO PRESCRIPTION MEDICATION SHOULD BE DISCONTINUED WITHOUT FIRST CONSULTING YOUR PHYSICIAN.**

